

↑  
GLOBOPHYCUS rugosum

Schopf 1968

Schopf 1968 p. 683-684 pl. 84 figs. 1a-c ; table 4

Division: Chlorophycophyta? Rhodophycophyta?  
Class: Chlorophyceae? Rhodophyceae?

- Vavrdová 2000 Neoproterozoic Czech Republic
- Qian, Yuan, Yan & Ding 2002 : Neoproterozoic, northern Jiangsu  
(p. 372 pl. 4, figs 6 & 9-10) & Anhui Provinces East China
- Schopf & Blacic 1971 : Late Precambrian Amadeus Basin  
(p. 110, fig. 13) Northern Territory Australia
- Schopf 1968 : Late Precambrian Central Australia
- Luo, Wang, Wang & Yin 1984 : latest Sinian / earliest Cambrian  
Guizhou Province China
- Sarkar 1989 (p. 33, pl. 2, fig. 3) Precambrian (1700-1350 My)  
Noamundi Basin, eastern India
- Kumar & Srivastava 1992 : Middle/Late Proterozoic (i.e. Rhyphen)  
(p. 304, Figs 9H) Garhwal Himalaya, India
- Qian, Wang, & Yan 2008 (p. 72 English p. 129, pl. 26, figs 6 & 9-11)  
Neoproterozoic SE margin of North China
- Kumar & Maithy 2008 : Ediacaran  
Krol Belt Lesser Himalaya India
- Maithy & Kumar 2007 : ditto
- Xiao, Hu, Ding & Liu & Ding et al. 1996 : Late Sinian  
eastern region of Yangtze Gorges, Zigui County, Hubei Province China