

MULTIPLICISPHAERIDIUM birminghamensis^{*} Cramer^{ex}, Eivenack et al 1973

1968 *Balloisphaeridium birminghamensis* Cramer, p. 127 nom nud

1970a *Balloisphaeridium birminghamensis* Cramer, p. 177-178, pl. 22, fig 314, 316 holotype not designated

1973 *Multiplicisphaeridium birminghamensis* (Cramer) Eivenack, Cramer & Diez, p. 539
holotype designated as pl 22, fig 314

* o.c. birminghamense

> HOESKLINTIA by (Sarjeant & Vavrdová 1997 p. 25)
Le Hérisse 1989 p. 146

- Cramer 1970a : late Llandoveryan to earliest Wenlockian, Alabama USA
- Duffield & Legault 1982 (pl. 1, fig. 10) : Llandoveryan, Quebec, Canada
- Diez & Cramer 1974 : late Llandoveryan - early Wenlockian range
- Cramer & Diez 1970 : Early Silurian USA/Canada
- Cramer 1968d : Early Silurian, Alabama USA
- Duffield & Legault 1981 (pl. 1, fig. 9) Early Silurian,
Anticosti Island Québec & Canada
- Cramer 1969f (pl. 6, fig. 3) : Ludlow, Belmonte, Asturias, N. Spain
- Cramer 1969b : Early Silurian, Alabama USA