

MULTIPLICISPHAERIDIUM <sup>ex</sup> bonitum : Cramer, Eisenack, Cramer & Diez 1973

p 541-542

1968 *Baltesphaeridium* sp 7 Gardine & Yapudjian, p. 463, pl. 3, fig. 11

1968 *Baltesphaeridium* sp Cramer, p 93, pl. 7, fig. 3

1970 *Baltesphaeridium bonitum* Cramer, p. 150-152, pl. 15, fig. 207-209

holotype not designated

⊗ 1970 *Thysanoprobolus polykion* Loeblich-Tappen, p. 262-266, fig. 1-12.

⊗ jr. syn. of *THYSANOPROBOLUS polykion* according to Eisenack et al 1973  
Who considered *M. bonitum* to be the senior name

> *VISBYSPHAERA* by Le Hérisse 2002 p. 390

- Cramer 1970a : Ludlovian - basal Early Gedinnian, Spain
- Gardiner & Yopandjian 1968 : late Nilusian, Libya
- Gardiner et al 1974 : Ludlovian, Algeria
- Diez & Cramer 1974 : Ludlovian - early Siegerian range
- Cramer Diez & Cuerdo 1974 : Early Ludlovian Bolivia
- Vavříková Isaacson et al. 1993 : Late Famennian, Bolivia
- Stemans, Le Hérisse & Bozdoğan 1996 : Early Prídolí, SE Turkey
- Hu 2001 (p. 69 pl. 3, fig. 1) : Arenig, Sandu Area, Guizhou Province  
China (biozone N<sub>2</sub>)
- Jachowicz 2000 (pl. 4, figs. 3, 29-10 & 12, pl. 5, figs. 10-11)  
Late Ludlow (Late Ludfordian)  
Pomeranian Caledonides, northern Poland