

NAVIFUSA *brasiliensis* *lingula* (Brito - Santos) Combar,

Lange & Panvart 1967^{*}
p 295

1965 *Leiofusa brasiliensis lingula* B.-S., p. 8, pl. 1, fig 1, pl. 2, fig. 2 invalid; taxonomic rank not indicated

1970a *N. brasiliensis lingula* (B.-S.) Loeblich, p. 731

* name not validly published

jr. syn. of *N. bacilla* according to Playford 1977 p. 29 and Wicander & Wood 1981 p 52

a Silurian (Zone T)

- Brito - Santos 1965 : Early Devonian? (Zone R), Brazil
- Kombaz et al 1967 : range Llandoveryan - Emurian; S. America
- Brito 1967a (p. 474-475) : Silurian, Early Devonian?, Maranhao Basin, Brazil
- Legault 1982 (pl. 1, fig. 2) : late Karadocian - Ashgillian, Labrador Sea
- Diez and Cramer 1977a : Gedinian - Givetian range uncertain Brazil
- Brito 1971 : Late Ordovician / Early Silurian - Early / Middle Devonian
- Bär & Riegel 1980 (Figure 3.8) : latest Ordovician / earliest Silurian Ghana
- Bär & Riegel 1974 (pl. 2 fig. 6) Frasnian Ghana
- Fang & He 1988 (pl. 1, fig. 26) : Silurian, Western Zhejiang China
- Xing & Fang 1999 (pl. 15, fig. 2) Llandovery / Wenlock Xinjiang China
- Wang, Gao Wang & Jiang 2008 (fig. 3.9) : Arenig / Llanvirn
Northern Qinling Orogenic Belt, Central China