

T

PERTICIELLA songliacensis

He, Gao & Qiao 1992 p. 188 pl. 2, figs 17-20 (holotype 1♂)
English translation p. ~~191~~ 195

He, Gao & Qiao 1992 : Albion NE China

Gao, Qiao & He 1992 : Albion NE China

Wang, Qin, Wu & Li 1996 (pl. 1, figs. 14-15) 'Campanian' Nei Monggol

(cf) Qiao et al. 2003: Hauterivian / Barremian,
Yangji Basin, Jilin, NE China

Perticiella songliaoensis gen. et sp. nov.

(Pl. I, figs. 17-20)

Description Shell cylindrical to somewhat dumbbell-shaped in outline, much longer than wide, straight or curved; both ends slightly expanded, rounded-square to rounded, with labial thickened free ends occupied by a circular central depression (about $4\ \mu\text{m}$ in diameter) not connecting with interior of shell. Wall thin, single-layered, light-colored; thickened free ends brown in color; wall surface ornamented with fine granules and short rugae. Opening rectangular $12\ \mu\text{m} \times 4.8\ \mu\text{m}$ in size, located at the centre of shell, with its long axis parallel to that of shell; operculum free.

Dimensions Shell $36-47\ \mu\text{m}$ long; middle area $9.6-14.2\ \mu\text{m}$ wide; free ends $12-14.2\ \mu\text{m}$ wide.

Occurrence Member 3 of Quantou Formation, Zhaozhou County, Heilongjiang.



