


Hemipodorhabdus latiforatus BLACK, 1971



6

Fig. 6 — *Hemipodorhabdus latiforatus* n. sp. Compound Nodular Bed, D1, Speeton. Holotype: H.710/25111. Distal view. x 6000.

Description:

A species of *Hemipodorhabdus* with two concentric cycles of plates on the distal surface, and D-shaped windows more than one micron in breadth.

Size: Distal shield $5.0 \times 4.1 \mu$, to $5.8 \times 4.6 \mu$, 24-27 rays; inner cycle $3.7 \times 2.4 \mu$ to $4.1 \times 2.7 \mu$; central opening $2.4 \times 1.1 \mu$ to $2.9 \times 1.2 \mu$. Specimens measured, 6.

Type level:

Hauterivian and Aptian.

Occurrence: Beds D1 and Upper B of the Speeton Clay at Speeton.

Type locality:

Bed D1 (Hauterivian) of the Speeton Clay at Speeton, England.

Depository:

Sedgwick Museum, Cambridge. Holotype: H.710/25111.

Author:

Black M., 1971, p. 404; pl. 31, fig. 6.

Reference:

Coccoliths of the Speeton Clay and Sutterby Marl. Proc. Yorks. Geol. Soc., vol. 38, part 3, n° 18, pp. 381-424, pls. 30-34, 3 tables.